

MATH 220 - Lecture 23 (11/05/2013)

23.1

Homework on Section 2.8 due Friday, Nov 8.

Section 2.9 due Tuesday, Nov 12.

Recall (results on bases of subspaces, dimension, rank, etc.)

Basis of a subspace H is an LI set of vectors in H that spans H .

dimension of H , denoted by $\dim H$ or $\dim(H)$, is the # vectors in any basis of H .

$\dim(\{\bar{0}\}) = 0$ as $\{\bar{0}\}$ is not LI.

$\text{rank}(A) = \dim(\text{Col } A) = \# \text{ pivot columns}$

$\dim(\text{Nul } A) = \# \text{ free variables}$

A basis for $\text{Col } A$: pivot columns of A

A basis for $\text{Nul } A$: vectors in parametric vector form of solutions to $A\bar{x} = \bar{0}$.

Rank theorem

For $A \in \mathbb{R}^{m \times n}$,

$$\text{rank}(A) + \dim(\text{Nul } A) = n$$

pivot columns

free variables

total # columns

Invertible matrix theorem (continued...)

- (a) $A \in \mathbb{R}^{n \times n}$ is invertible.
- (m) Columns of A form a basis for \mathbb{R}^n .
- (n) $\text{Col } A = \mathbb{R}^n$.
- (o) $\dim \text{Col } A = n$.
- (p) $\text{rank } A = n$.
- (q) $\text{Nul } A = \{\vec{0}\}$.
- (r) $\dim(\text{Nul } A) = 0$.

We could use any basis to represent a subspace. In any given basis, we can talk about the "coordinates" of any vector in the subspace.

Coordinates

Let $B = \{\vec{b}_1, \dots, \vec{b}_p\}$ be a basis for a subspace H . Each $\vec{x} \in H$ can be written as $\vec{x} = c_1 \vec{b}_1 + c_2 \vec{b}_2 + \dots + c_p \vec{b}_p$ for scalars c_1, c_2, \dots, c_p , which are unique to \vec{x} . These scalars are called the coordinates of \vec{x} relative to the basis B . Stacking these scalars into a vector

$$[\vec{x}]_B = \begin{bmatrix} c_1 \\ c_2 \\ \vdots \\ c_p \end{bmatrix} \text{ is the } B\text{-coordinate of } \vec{x}.$$

When B is the standard basis, i.e., $\{\vec{e}_1, \dots, \vec{e}_p\}$, the B -coordinate of any \vec{x} consists of its own entries.
(So, the B -coordinate of \vec{x} is \vec{x} itself here)

Pg 157.

In Exercises 1 and 2, find the vector \mathbf{x} determined by the given coordinate vector $[\mathbf{x}]_{\mathcal{B}}$ and the given basis \mathcal{B} . Illustrate your answer with a figure, as in the solution of Practice Problem 2.

$$1. \mathcal{B} = \left\{ \begin{bmatrix} 1 \\ 1 \end{bmatrix}, \begin{bmatrix} 2 \\ -1 \end{bmatrix} \right\}, [\mathbf{x}]_{\mathcal{B}} = \begin{bmatrix} 3 \\ 2 \end{bmatrix}$$

$$\bar{\mathbf{x}} = 3 \begin{bmatrix} 1 \\ 1 \end{bmatrix} + 2 \begin{bmatrix} 2 \\ -1 \end{bmatrix} = \begin{bmatrix} 7 \\ 1 \end{bmatrix}$$

→ We just have to evaluate the linear combination given

In Exercises 3–6, the vector \mathbf{x} is in a subspace H with a basis $\mathcal{B} = \{\mathbf{b}_1, \mathbf{b}_2\}$. Find the \mathcal{B} -coordinate vector of \mathbf{x} .

$$5. \mathbf{b}_1 = \begin{bmatrix} 1 \\ 4 \\ -3 \end{bmatrix}, \mathbf{b}_2 = \begin{bmatrix} -2 \\ -7 \\ 5 \end{bmatrix}, \mathbf{x} = \begin{bmatrix} 2 \\ 9 \\ -7 \end{bmatrix}$$

$\mathcal{B} = \{\bar{\mathbf{b}}_1, \bar{\mathbf{b}}_2\}$. Find $[\bar{\mathbf{x}}]_{\mathcal{B}}$. So, find c_1, c_2 such that

$$\bar{\mathbf{x}} = c_1 \bar{\mathbf{b}}_1 + c_2 \bar{\mathbf{b}}_2$$

$$\begin{array}{c} \textcircled{1} \\ \left[\begin{array}{cc|c} -2 & 2 & 2 \\ 4 & -7 & 9 \\ -3 & 5 & 7 \end{array} \right] \xrightarrow[\substack{R_2 - 4R_1 \\ R_3 + 3R_1}]{R_2 - 4R_1} \left[\begin{array}{cc|c} 1 & -2 & 2 \\ 0 & \textcircled{1} & 1 \\ 0 & -1 & -1 \end{array} \right] \xrightarrow[\substack{R_1 + 2R_2 \\ R_3 + R_2}]{R_3 + R_2} \left[\begin{array}{cc|c} 1 & 0 & 4 \\ 0 & 1 & 1 \\ 0 & 0 & 0 \end{array} \right] \end{array}$$

$$c_1 = 4, \\ c_2 = 1$$

$$\text{or } [\bar{\mathbf{x}}]_{\mathcal{B}} = \begin{bmatrix} 4 \\ 1 \end{bmatrix}$$

Why study coordinates?

One often works with a nonstandard basis for a subspace. Hence we want to study how any vector is expressed in such a basis. For instance, we could start with the standard basis for an image, and do a bunch of geometric transformations, e.g., rotate 90° ccw. After that step, we could just work with the nonstandard basis for further analyses.

Pg 158.

In Exercises 17 and 18, mark each statement True or False. Justify each answer. Here A is an $m \times n$ matrix.

17. a. If $\mathcal{B} = \{\mathbf{v}_1, \dots, \mathbf{v}_p\}$ is a basis for a subspace H and if $\mathbf{x} = c_1\mathbf{v}_1 + \dots + c_p\mathbf{v}_p$, then c_1, \dots, c_p are the coordinates of \mathbf{x} relative to the basis \mathcal{B} .
- b. Each line in \mathbb{R}^n is a one-dimensional subspace of \mathbb{R}^n .
- c. The dimension of $\text{Col } A$ is the number of pivot columns in A .
- d. The dimensions of $\text{Col } A$ and $\text{Nul } A$ add up to the number of columns in A .
- e. If a set of p vectors spans a p -dimensional subspace H of \mathbb{R}^n , then these vectors form a basis for H .

- (a). True. Definition of coordinates.
- (b) False. Only if it goes through the origin.
- (c) True. Definition.
- (d) True. Rank theorem
- (e) True. A set of p vectors that spans a p -dimensional subspace will be LI. Hence it's a basis.

Pg 159

23. If possible, construct a 3×5 matrix A such that $\dim \text{Nul } A = 3$ and $\dim \text{Col } A = 2$.

$$n=5, \quad \dim \text{Col } A + \dim \text{Nul } A = 2 + 3 = n \quad \checkmark$$

(So, rank theorem holds)

$$A = \begin{bmatrix} 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix} \quad \text{works. (has 2 pivot columns).}$$

Determinants (Section 3.1)

Recall, $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$, then $A^{-1} = \frac{1}{ad-bc} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix}$, when

determinant of A , $\det A = ad-bc \neq 0$.

In general, for $A \in \mathbb{R}^{n \times n}$, A is invertible if and only if $\det A \neq 0$.

Let $A = \begin{bmatrix} 1 & 2 & 3 \\ 4 & a & b \\ 5 & c & d \end{bmatrix}$. When is A invertible?
(What condition should a, b, c, d satisfy in order to make A invertible?)

$$\begin{bmatrix} 1 & 2 & 3 \\ 4 & a & b \\ 5 & c & d \end{bmatrix} \xrightarrow{\substack{R_2 - 4R_1 \\ R_3 - 5R_1}} \begin{bmatrix} 1 & 2 & 3 \\ 0 & a-8 & b-12 \\ 0 & c-10 & d-15 \end{bmatrix} \xrightarrow{R_3 - \left(\frac{c-10}{a-8}\right)R_2} \rightarrow$$

$$\begin{bmatrix} 1 & 2 & 3 \\ 0 & a-8 & b-12 \\ 0 & 0 & (d-15) - \left(\frac{c-10}{a-8}\right)(b-12) \end{bmatrix} \neq 0$$

as we need three pivots

$$(a-8)(d-15) - (c-10)(b-12) \neq 0$$

$$(ad - 15a - 8d + 120) - (bc - 12c - 10b + 120) \neq 0$$

$$(ad - bc) - (8d - 10b) + (12c - 15a) \neq 0$$

i.e., we need $1(ad-bc) - 2(4d-5b) + 3(4c-5a) \neq 0$
 \rightarrow expanding along Row 1

$$\det \begin{pmatrix} \boxed{1 \ 2 \ 3} \\ 4 \ a \ b \\ 5 \ c \ d \end{pmatrix} = 1 \cdot \det \begin{pmatrix} a & b \\ c & d \end{pmatrix} - 2 \det \begin{pmatrix} 4 & b \\ 5 & d \end{pmatrix} + 3 \det \begin{pmatrix} 4 & c \\ 5 & a \end{pmatrix}$$

We will see that one could expand along any row or any column to compute $\det A$. More on the details in the next lecture.